



**Territorial defense in a network
audiences only matter to male fiddler crabs primed for confrontation**

Darden, Safi K.; May, Maggie K.; Boyland, Natasha K.; Dabelsteen, Torben

Published in:
Behavioral Ecology

DOI:
[10.1093/beheco/ary169](https://doi.org/10.1093/beheco/ary169)

Publication date:
2019

Document version
Peer reviewed version

Citation for published version (APA):
Darden, S. K., May, M. K., Boyland, N. K., & Dabelsteen, T. (2019). Territorial defense in a network: audiences only matter to male fiddler crabs primed for confrontation. *Behavioral Ecology*, 30(2), 336-340.
<https://doi.org/10.1093/beheco/ary169>

1 Territorial defence in a network: audiences only matter to male fiddler crabs primed for
2 confrontation

3
4 **Lay Summary:** Being part of a social network means that responses to social confrontations
5 are likely to be more complex than they might seem. Indeed, here we find effects of a wider
6 network of conspecifics on an individual's behaviour in male European fiddler crabs. Males
7 became more aggressive toward intruders if their neighbour was watching when they had
8 previously observed an aggressive interaction between their neighbour and a male territory
9 intruder.

10 11 **Abstract**

12
13 Territorial contests often occur in the presence of conspecifics not directly involved in the
14 interaction. Actors may alter their behaviour in the presence of this audience, an 'audience
15 effect', and audiences themselves may alter their behaviour as a result of observing an
16 interaction, a 'bystander effect'. Previous work has documented these effects by looking at
17 each in isolation, but to our knowledge, none has investigated their interaction; something
18 that is more likely to represent a realistic scenario for species where individuals aggregate
19 spatially. We therefore have a somewhat limited understanding of the extent and direction of
20 these potentially complex indirect social effects on behaviour. Here we examined how
21 audience and bystander effects work in tandem to modify resident male aggressive behaviour
22 towards intruders in European fiddler crabs, *Afruca tangeri*. We found that male crabs with
23 an audience showed greater aggressive behaviour towards an intruder compared to males
24 without an audience, but only if they had acted as a bystander to an aggressive signalling
25 interaction prior to the intrusion. Indeed, bystanding during aggressive interactions elevated

aggressive responses to intruders maximally if there was an audience present. Our results suggest that bystanding had a priming effect on territory-holding males, potentially by providing information on the immediate level of competition in the local neighbourhood, and that same-sex audiences only matter if males have been primed. This study highlights the fundamental importance of considering broader interaction networks in studying real-world dyadic interactions and of including non-vertebrate taxonomic groups in these studies.

Key words: audience effect, bystander effect, invertebrate, fiddler crab, *Afruca tangeri*, *Uca tangeri*

Background

Considering communication in the context of a wider network has revealed behaviours and effects such as eavesdropping, audience effects and bystander effects, that would not be observable from a dyadic approach (McGregor, 2005). Territorial contests are typically settled through pair-wise interactions within a network of multiple individuals, rather than in (McGregor and Dabelsteen, 1996). Audience effects occur when animals modify their behaviour due to the presence of other individuals not involved in the interaction (Zuberbühler, 2008) and these effects have been shown to alter the intensity of agonistic displays in a number of species (Cruz and Oliveira, 2015; dos Santos et al., 2017; Fitzsimmons and Bertram, 2013; Montroy et al., 2016; Setoguchi et al., 2015). In the context of territorial defence, the audience effect has been demonstrated to be dependent not only on the sex of the audience, but also on the territorial status and familiarity of individuals in the wider network (Bertucci et al., 2014; Dzieweczynski et al., 2005). Observed differences in behaviour in the presence of an audience suggest that individuals can (1) assess attributes of their audience and (2) adjust their behaviour as a strategy to counter costs (or strengthen benefits) that can come with eavesdropping, or more generally, gathering of social information by audiences (Earley and Dugatkin, 2002). In the bystander effect the audience members themselves are influenced by observing an interaction (Earley and Dugatkin, 2002; Oliveira et al., 2001; Peake et al., 2006). The observation acts directly on the motivational system (Hirschenhauser and Oliveira, 2006; Oliveira et al., 2001) and prepares individuals for what may happen next in their social environment (Antunes and Oliveira, 2009). Individuals can, for example, be primed to augment their levels of aggression in interactions following bystanding. For example, Clotfelter and Paolino (Clotfelter and Paolino, 2003) found increased aggressiveness by Siamese fighting fish, *Betta splendens*, towards a novel male

conspecific after the observation of an aggressive interaction. However, reverse priming, a reduction in aggressive behaviour, has also been found, for example, in a study of crayfish, *Orconectes rusticus* (Zulandt et al., 2008), which to our knowledge is also the only invertebrate species in which bystander effects have been investigated.

Although audience and bystander effects are frequently documented, it is not clear from the literature how bystander and audience effects may interact. This represents a potential key gap in our understanding of the strength or importance of these effects in *in situ* contexts where they are likely to happen concurrently, particularly in species where individuals live in aggregated communities. A study in Siamese fighting fish, suggests that simply viewing an unfamiliar male prior to a contest with another male can prime males for increased aggression during a fight, irrespective of whether the audience is present or not during the fight itself (Matos et al., 2003). It also highlights the complexity of bystander and audience effects and the need to manipulate these effects in tandem in order to deepen our understanding of the significance of dynamics in the social environment for the expression of behaviour. Here we investigate the interplay between audience and bystander effects on the territorial behaviour of male European fiddler crabs, *Afruca tangeri*. Fiddler crabs are very unlikely to be found in isolation (Pope, 2005), and therefore have a high likelihood of engaging in social interactions with multiple receivers within signal range. Males actively defend territories around their burrows (Hemmi and Zeil, 2003) with their one greatly enlarged claw, used for the production of highly conspicuous visual agonistic and courtship signals and for fighting (Oliveira and Custódio, 1998; Wolfrath, 1993). Male crabs without a burrow will wander through the population and challenge burrow holders in attempts to acquire a burrow (Jordao and Oliveira, 2005). Burrow holding males must therefore invest in an agonistic response in order to retain residency (Oliveira and Custódio, 1998). Recent work in another fiddler crab

species suggests that same-sex audiences do not matter for expressed levels of aggression (dos Santos et al., 2017), given the ubiquity of males in the immediate social environment (i.e. males are highly likely to have male neighbours), this is perhaps not surprising. We propose instead that males will fine-tune their responsiveness to their social environment according to current social information, such as the immediate level of competition in the neighbourhood. If males observe a territorial dispute in their neighbour's territory, they should be primed for a possible challenge of their own territory and therefore will be more responsive or sensitive to their immediate social environment. In this case, bystanding to an aggressive interaction between a neighbouring male and a stranger (intruding male), should augment the aggressive response exhibited by males towards intruders appearing close in time. This should be particularly so when the social stimuli are increased by a male audience as this audience can gather information on the interaction outcome (e.g., the competitive ability of his male neighbour) and in the least, acts as an indicator of a more competitive social environment (i.e. more males are present) than when no male audience is present. In this investigation, we used an *in situ* experimental manipulation of the social environment that burrow holding male crabs experienced to test such effects during territorial confrontations in the home environment.

Methods

The study was carried out in the Parque Natural da Ria Formosa, Portugal (N370927, E073244) from May to July 2012 in order to quantify audience and bystander effects *in situ* in males of a free roaming population of European fiddler crabs. To test focal male crabs at their home burrow an arena made of bamboo and sand-coloured fabric was placed around two neighbouring males and their burrows (Fig. 1). Neighbouring male pairs were selected

according to three criteria: (1) they were matched in claw size (estimated by visual comparison); (2) they were exhibiting courtship behaviour (courtship waving) indicating that they were actively defending a burrow; and (3) they were within one meter of one another (measured using a measuring tape stretched in a straight line from the center of one burrow to the other). Stimulus crabs captured from other areas of the mudflat were tethered with 10cm of clear monofilament line to bamboo posts inserted into the substrate within the arena to simulate wandering male crabs (intruders). This method of tethering stimulus crabs is relatively standard and has been used successfully in this and several other fiddler crab species to elicit both courtship and territorial behaviour as observed under natural conditions (Detto and Backwell, 2009; Detto et al., 2006; Detto et al., 2010; How et al., 2008; Milner et al., 2010; Pope, 2005; Reaney, 2007)(Bookmythe et al., 2010). Intruders were matched in claw size to focal crabs using visual comparison. Focal crabs were exposed to two 5-minute phases: a ‘bystanding phase’ (neighbour interacts with a simulated intruder) and a subsequent ‘interaction phase’ (focal male interacts with a simulated intruder) in one of four treatments (3 control and 1 experimental): (1) *null control* – no neighbour-intruder interaction in the ‘bystanding phase’ and no audience in the ‘interaction phase’ (n=11); (2) *audience control* – no neighbour-intruder interaction in the ‘bystanding phase’ and an audience in the ‘interaction phase’ (n=10); (3) *bystander control* – neighbour-intruder interaction in the ‘bystanding phase’ and no audience in the ‘interaction phase’ (n=11); and (4) *bystander and audience (experimental)* – neighbour-intruder interaction in the ‘bystanding phase’ and audience in the ‘interaction phase’ (n=11) (Fig. 1).

In the ‘bystanding phase’ of treatments (3) and (4) a stimulus male was tethered 40cm from the neighbour’s burrow (min 90 cm from the focal male’s burrow) and left for 5 minutes after both males (focal and neighbour) had emerged from their burrows. In the ‘bystanding phase’

of treatments (1) and (2) the arena was approached (and stimulus male placement simulated) and the crabs were then left for 5 minutes after both males had emerged from their burrow. In the interaction phase a stimulus male was tethered 40cm from the focal male's burrow to simulate a wandering male. In treatments (2) and (4) the neighbouring male was allowed to emerge from his burrow and act as an audience while in treatments (1) and (3) the neighbouring male's burrow was blocked to prevent him from emerging during the trial period. Phases began when the focal or both males had surfaced (carapace and major claw visible), as applicable, and lasted for five minutes. Following a trial, burrows were marked with a small flag and within a semi-lunar tidal cycle these areas were avoided for further testing to ensure that males were not re-used over the course of the study. We quantified aggression as a behavioural state by measuring the duration of time that focal males were engaged in aggressive behaviours toward the simulated intruder in the 5-minute interaction phase. The observed aggressive behaviour, following previously published methods (see Burford et al., 2000; Oliveira et al., 1998; Wolfrath, 1993), included both non-contact aggression (threat displays) and that involving physical contact (pushing, grappling and tossing) and the intensity of enactment represents the focal male's willingness to escalate the contest (Oliveira et al., 1998; Wolfrath, 1993). We compared the time focal males spent performing aggressive behaviour among treatments with an analysis of variance with treatment as a fixed effect (SPSS v. 22). We included inter-burrow distance between the focal and neighbour as a covariate in the model as this varied across focal individuals. Post hoc analyses were carried out on the marginal means using a least significant difference adjustment for multiple comparisons. All behaviours were scored from video recordings (Panasonic HDC-SD800 camcorder) of the trials by a single observer (MKM) naïve to the trial condition.

The study was carried out under permit (ICNF, Portugal) following ethical approval from the first author's home institution. All captured crabs were kept singly in shaded containers filled with sea water and mud prior to testing and released back to their area of capture at the end of a trial. Crabs were marked with non-toxic paint to ensure that they were not used more than once in case of recapture.

Results

There was an overall effect of the treatment that focal males experienced on the amount of aggressive behaviour they displayed toward a simulated intruder ($F_{3,38}=11.797$, $p<0.0001$, Fig. 2). Our post hoc analysis revealed that there was an effect of having a neighbouring male audience present during a territorial intrusion on the level of aggression expressed by a burrow-holding male, but only if burrow-holding males had previously acted as bystander (Treatment 4) to a neighbouring aggressive interaction (Table 1; Fig. 2). In short, males in Treatment 4 behaved more aggressively than in any other treatment. Males in two of the control treatments (Treatments 1 and 2) did not differ from one another in behaviour, but males in the null control (no bystanding and no audience; Treatment 1), were less aggressive than males in the bystander control (bystanding, no audience) (Treatment 3).

Discussion

Gleaning information from the social environment is essential for an individual to respond to this environment appropriately (Danchin et al., 2004; Seppanen et al., 2007; Valone, 2007). Individuals living in systems that are likely to be high in social noise will be under particular pressure to either filter out information to avoid an inappropriate response or to use specific

cues to alert them to a situation where an active response may be required (e.g., Fitzsimmons et al., 2008; Naguib et al., 2004). This latter priming effect may be particularly important in predicting that, for example, a territorial challenge is likely to occur. The anticipation of being challenged is likely to affect not only how individuals respond to a rival conspecific during a direct interaction, but also how they respond to the presence of other conspecifics in the immediate environment. These individuals that are not part of the interaction, may themselves be gleaned social information (i.e. they are conspecific audiences). In this study we found the first evidence that a neighbouring male audience moderated male European fiddler crab behaviour towards an intruder, but only if males had acted as bystander to an aggressive interaction between a male neighbour and a same-sex intruder immediately prior to the encounter. This suggests that focal males that had viewed an aggressive interaction between his neighbour and an intruder were somehow primed for heightened responsiveness to having a male audience present during an interaction between himself and a territorial intruder. The observed increase in aggressiveness was likely also a function of a general priming for an aggressive response as a result of the bystanding, since we did detect greater aggressiveness when males without an audience had acted as bystander compared to males in our null control (no audience and no bystanding).

The prevalence of the use of public information (Danchin et al., 2004) suggests that it could be beneficial for individuals to employ strategies to manage the perceptions of unintended receivers. Fighting could communicate the focal male's motivation to defend his territory (Detto et al., 2010) or ability to do so (Peake et al., 2001) to his neighbour. However, we did not see an effect of a neighbouring male audience unless the focal male had previously acted as a bystander to an aggressive interaction between his neighbour and an intruder. Bio-regulatory mediators of the adjustment of aggressive behaviour are likely to be affected in

males that anticipate a territorial challenge based on cues in their social environment; facilitating an adjustment to an increased competitive environment with the appropriate behavioural response (Antunes and Oliveira, 2009). It could be that the simple presence of a neighbour is not enough of a cue of the extent of the competitive environment, given the ubiquity of territorial neighbours, and their familiarity, in a male's visual field (Detto et al., 2010; Hemmi and Zeil, 2003; Pope, 2005). Instead, if a territorial challenge has been observed immediately prior to an intrusion, effects on bio-regulatory mechanisms (Oliveira et al., 2001) could drive a response directly or at least provide a cue as to the probability of an escalation of aggression that could lead to a territorial takeover (Oliveira, 2009).

The bystander effect detected in this study is consistent with work in vertebrate species demonstrating increased levels of aggression (e.g., Clotfelter and Paolino, 2003), but inconsistent with the reverse priming demonstrated in another crustacean species (Zulandt et al., 2008). In our study, burrow holding males will have invested energy into establishing territory boundaries with their neighbours (Detto et al., 2010) and as such intruders are a potential threat to the territory more generally if they were to take over a neighbour's burrow. Zulandt and colleagues (Zulandt et al., 2008) suggest that the bystander effect could be resource dependent such that when resources are abundant, and consequently their value reduced, an individual will avoid escalation if the observation of a fight indicates that there is an increased likelihood of being challenged within their network. In the present study males were at their home burrow and as a consequence were defending a valuable resource; an added ecological realism that may have contributed to the direction of the effect.

Our study demonstrates that there is likely to be a complex interaction between audience and bystander effects in systems where individuals are part of a network of conspecific

interactants. The results suggest that selection has enhanced the use of public information to increase the appropriateness of responses to social stimuli. We investigated effects of the presence of a fiddler crab's nearest neighbour, an individual that the focal is likely to have interacted with before and to be familiar with, however, laboratory work on audience effects has demonstrated that attributes of an audience can influence physiological and behavioural effects on individuals, for example, sex and territorial status (Dzieweczynski et al., 2005). Future *in situ* work investigating behavioural reactions when the audience is a *non*-neighbour or a female, would therefore be particularly enlightening.

References

- Antunes RA, Oliveira RF, 2009. Hormonal anticipation of territorial challenges in cichlid fish. *Proceedings of the National Academy of Sciences* 106:15985-15989. doi: 10.1073/pnas.0900817106.
- Bertucci F, Matos RJ, Dabelsteen T, 2014. Knowing your audience affects male–male interactions in siamese fighting fish (*Betta splendens*). *Animal Cognition* 17:229-236. doi: 10.1007/s10071-013-0655-6.
- Booksmythe I, Jennions MD, Backwell PRY, 2010. Interspecific assistance: Fiddler crabs help heterospecific neighbours in territory defence. *Biology Letters* 6:748-750. doi: 10.1098/rsbl.2010.0454.
- Burford FRL, McGregor PK, Oliveira RF, 2000. Response of fiddler crabs (*Uca tangeri*) to video playback in the field. *Acta Ethologica* 3:55-59. doi: 10.1007/s102110000020.
- Clotfelter ED, Paolino AD, 2003. Bystanders to contests between conspecifics are primed for increased aggression in male fighting fish. *Animal Behaviour* 66:343-347. doi: <http://dx.doi.org/10.1006/anbe.2003.2227>.

260 Cruz AS, Oliveira RF, 2015. Audience effects and aggressive priming in agonistic behaviour
 261 of male zebrafish, *Danio rerio*. *Animal Behaviour* 107:269-276. doi:
 262 <http://dx.doi.org/10.1016/j.anbehav.2015.07.007>.

263 Danchin É, Giraldeau L-A, Valone TJ, Wagner RH, 2004. Public information: From nosy
 264 neighbors to cultural evolution. *Science* 305:487-491. doi: 10.1126/science.1098254.

265 Detto T, Backwell PRY, 2009. The fiddler crab *Uca mjoebergi* uses ultraviolet cues in mate
 266 choice but not aggressive interactions. *Animal Behaviour* 78:407-411. doi:
 267 [10.1016/j.anbehav.2009.05.014](http://dx.doi.org/10.1016/j.anbehav.2009.05.014).

268 Detto T, Backwell PRY, Hemmi JM, Zeil J, 2006. Visually mediated species and neighbour
 269 recognition in fiddler crabs (*Uca mjoebergi* and *Uca capricornis*). *Proceedings of the*
 270 *Royal Society B-Biological Sciences* 273:1661-1666. doi: [10.1098/rspb.2006.3503](http://dx.doi.org/10.1098/rspb.2006.3503).

271 Detto T, Jennions MD, Backwell PRY, 2010. When and why do territorial coalitions occur?
 272 Experimental evidence from a fiddler crab. *The American Naturalist* 175:E119-E125.
 273 doi: [10.1086/651588](http://dx.doi.org/10.1086/651588).

274 dos Santos LC, de Moraes Freire FA, Luchiari AC, 2017. The effect of audience on
 275 intrasexual interaction in the male fiddler crab, *Uca maracoani* (latreille 1802–1803,
 276 decapoda, ocypodidae). *Journal of Ethology* 35:93-100. doi: [10.1007/s10164-016-](http://dx.doi.org/10.1007/s10164-016-0495-3)
 277 [0495-3](http://dx.doi.org/10.1007/s10164-016-0495-3).

278 Dzieweczynski TL, Earley RL, Green TM, Rowland WJ, 2005. Audience effect is context
 279 dependent in siamese fighting fish, *Betta splendens*. *Behavioral Ecology* 16:1025-
 280 1030. doi: [10.1093/beheco/ari088](http://dx.doi.org/10.1093/beheco/ari088).

281 Earley RL, Dugatkin LA, 2002. Eavesdropping on visual cues in green swordtail
 282 (*Xiphophorus helleri*) fights: A case for networking. *Proceedings of the Royal Society*
 283 *B: Biological Sciences* 269:943-952. doi: [10.1098/rspb.2002.1973](http://dx.doi.org/10.1098/rspb.2002.1973).

284 Fitzsimmons LP, Bertram SM, 2013. Playing to an audience: The social environment
 285 influences aggression and victory displays. *Biology Letters* 9:20130449. doi:
 286 10.1098/rsbl.2013.0449.

287 Fitzsimmons LP, Foote JR, Ratcliffe LM, Mennill DJ, 2008. Eavesdropping and
 288 communication networks revealed through playback and an acoustic location system.
 289 *Behavioral Ecology* 19:824-829. doi: 10.1093/beheco/arn036.

290 Hemmi JM, Zeil J, 2003. Burrow surveillance in fiddler crabs - i. Description of behaviour.
 291 *Journal of Experimental Biology* 206:3935-3950. doi: 10.1242/jeb.00632.

292 Hirschenhauser K, Oliveira RF, 2006. Social modulation of androgens in male vertebrates:
 293 Meta-analyses of the challenge hypothesis. *Animal Behaviour* 71:265-277. doi:
 294 10.1016/j.anbehav.2005.04.014.

295 How MJ, Hemmi JM, Zeil J, Peters R, 2008. Claw waving display changes with receiver
 296 distance in fiddler crabs, *Uca perplexa*. *Animal Behaviour* 75:1015-1022. doi:
 297 10.1016/j.anbehav.2007.09.004.

298 Jordao JM, Oliveira RF, 2005. Wandering in male fiddler crabs (*Uca tangeri*): Alternative
 299 reproductive tactic or a functional constraint? *Behaviour* 142:929-939. doi:
 300 10.1163/1568539055010101.

301 Matos RJ, Peake TM, McGregor PK, 2003. Timing of presentation of an audience:
 302 Aggressive priming and audience effects in male displays of siamese fighting fish
 303 (*Betta splendens*). *Behavioural Processes* 63:53-61. doi:
 304 [https://doi.org/10.1016/S0376-6357\(03\)00029-9](https://doi.org/10.1016/S0376-6357(03)00029-9).

305 McGregor PK, editor 2005. Animal communication networks. Cambridge: Cambridge
 306 University Press.

307 McGregor PK, Dabelsteen T, 1996. Communication networks. In: Kroodsma DE, Miller EH,
 308 editors. *Ecology and evolution of acoustic communication* Ithaca, N.Y.: Cornell
 309 University Press. p. 409–425.

310 Milner RNC, Jennions MD, Backwell PRY, 2010. Safe sex: Male–female coalitions and pre-
 311 copulatory mate-guarding in a fiddler crab. *Biology Letters* 6:180-182. doi:
 312 10.1098/rsbl.2009.0767.

313 Montroy K, Loranger MJ, Bertram SM, 2016. Male crickets adjust their aggressive behavior
 314 when a female is present. *Behavioural Processes* 124:108-114. doi:
 315 <http://dx.doi.org/10.1016/j.beproc.2015.11.003>.

316 Naguib M, Amrhein V, Kunc HP, 2004. Effects of territorial intrusions on eavesdropping
 317 neighbors: Communication networks in nightingales. *Behavioral Ecology* 15:1011-
 318 1015. doi: 10.1093/beheco/arh108.

319 Oliveira RF, 2009. Social behavior in context: Hormonal modulation of behavioral plasticity
 320 and social competence. *Integrative and Comparative Biology* 49:423-440. doi:
 321 10.1093/icb/icp055.

322 Oliveira RF, Custódio MR, 1998. Claw size, waving display and female choice in the
 323 European fiddler crab, *Uca tangeri*. *Ethology Ecology & Evolution* 10:241-251. doi:
 324 10.1080/08927014.1998.9522855.

325 Oliveira RF, Lopes M, Carneiro LA, Canario AVM, 2001. Watching fights raises fish
 326 hormone levels. *Nature* 409:475-475.

327 Oliveira RF, McGregor PK, Burford FRL, Custodio MR, Latruffe C, 1998. Functions of
 328 mudballing behaviour in the European fiddler crab *Uca tangeri*. *Animal Behaviour*
 329 55:1299-1309.

330 Peake TM, Matos RJ, McGregor PK, 2006. Effects of manipulated aggressive ‘interactions’
 331 on bystanding male fighting fish, *Betta splendens*. *Animal Behaviour* 72:1013-1020.
 332 doi: <http://dx.doi.org/10.1016/j.anbehav.2006.01.029>.
 333 Peake TM, Terry AMR, McGregor PK, Dabelsteen T, 2001. Male great tits eavesdrop on
 334 simulated male-to-male vocal interactions. *Proceedings of the Royal Society B:*
 335 *Biological Sciences* 268:1183-1187.
 336 Pope DS, 2005. Waving in a crowd: Fiddler crabs signal in networks. In: McGregor PK,
 337 editor. *Animal communication networks* Cambridge: Cambridge University Press. p.
 338 252-276.
 339 Reaney LT, 2007. Foraging and mating opportunities influence refuge use in the fiddler crab,
 340 *Uca mjoebergi*. *Animal Behaviour* 73:711-716. doi: 10.1016/j.anbehav.2006.05.022.
 341 Seppanen JT, Forsman JT, Monkkonen M, Thomson RL, 2007. Social information use is a
 342 process across time, space, and ecology, reaching heterospecifics. *Ecology* 88:1622-
 343 1633.
 344 Setoguchi S, Kudo A, Takanashi T, Ishikawa Y, Matsuo T, 2015. Social context-dependent
 345 modification of courtship behaviour in *Drosophila prolongata*. *Proceedings of the*
 346 *Royal Society B: Biological Sciences* 282.
 347 Valone TJ, 2007. From eavesdropping on performance to copying the behavior of others: A
 348 review of public information use. *Behavioral Ecology and Sociobiology* 62:1-14. doi:
 349 10.1007/s00265-007-0439-6.
 350 Wolfrath B, 1993. Observations on the behaviour of the european fiddler crab *Uca tangeri*.
 351 *Marine Ecology-Progress Series* 100:111-118.
 352 Zuberbühler K, 2008. Audience effects. *Current Biology* 18:R189-R190. doi:
 353 <http://dx.doi.org/10.1016/j.cub.2007.12.041>.

354 Zulandt T, Zulandt-Schneider RA, Moore PA, 2008. Observing agonistic interactions alters
355 subsequent fighting dynamics in the crayfish, *Orconectes rusticus*. *Animal Behaviour*
356 75:13-20. doi: 10.1016/j.anbehav.2007.04.017.

357 Figure legends

358

359 Figure 1. Overview of experimental arenas and 4 treatments (3 control and 1 experimental)
360 used to investigate bystander and audience effects in male European fiddler crabs (see text for
361 details).


362

363 Figure 2. Average time that male European fiddler crabs spent engaging in aggressive
364 behaviour toward experimentally introduced male intruders in 4 social treatments (see Fig. 1)
365 designed to test for bystander and audience effects (Estimated Marginal Mean (EMM) \pm SE;
366 matching letters indicate treatments where $p > 0.05$ for tested differences).

367

Table 1. Results of the *post hoc* analysis of differences among treatments in burrow-holding male aggressive responses to simulated intrusions by conspecific male fiddler crabs. P-values in bold indicate where significant differences in aggressiveness were found.

Treatment A	Treatment B	Mean difference	95% CI for Difference		<i>P</i>
			Lower bound	Upper bound	
1 - Null control	2 - Audience control	-31.04	-82.41	20.33	0.229
	3 - Bystander	-53.73	-103.39	-4.07	0.035
	4 - Bystander and audience	-143.18	-194.50	-91.85	p<0.001
2 - Audience control	3 - Bystander	-22.69	-73.91	28.53	0.375
	4 - Bystander and audience	-112.14	-163.35	-60.92	p<0.001
3 - Bystander	4 - Bystander and audience	-89.45	-140.48	-38.42	0.001



375
376

